

Nulojix (belatacept)

Provider Order Form rev. 1/2/2026



PATIENT INFORMATION

Referral Status: New Referral Updated Order Order Renewal

Patient Full Name: _____ DOB: _____ Phone: _____ Gender: M F Other

Email Address: _____ Address: _____ Weight (lbs/kg): _____ Height (in): _____

NKDA Allergies: _____ Existing prior authorization? Yes, (Send a copy) No (AIC will process)

Patient Status: New to Therapy Continuing Therapy Last Treatment Date: _____ Next Due Date: _____

Patient Preferred Location: _____

DIAGNOSIS & CLINICAL INFORMATION

ICD 10-Code & Description (Provide full completed code)

Z94.0 Kidney transplant status
 Z48.22 Encounter for aftercare following kidney transplant

D82.3 Immunodeficiency following hereditary defective response to Epstein-Barr virus
 Other: _____

REQUIRED DOCUMENTATION: Please include insurance card (front & back), all patient demographics, history & physicals, medication lists, recent lab results (CBC, CMP, TB, Hep B panel-depending on medication) , signed prescription order and recent visit notes.

Confirm that these and the required lab orders have been sent to American Infusion Care and necessary parties.

PRESCRIPTION INFORMATION

Nursing: Provide nursing care per American Infusion Care - Specialty Infusions protocols, including reaction management and post-procedure observation

Pre-Medications

Acetaminophen (Tylenol) 500mg 650mg 1000mgPO
 Cetirizine (Zyrtec) 10mgPO
 Loratadine (Claritin) 10mgPO
 Diphenhydramine (Benadryl) 25mg 50mg PO IV
 Methylprednisolone (Solu-Medrol) 40mg 125mg IV
 Other: _____ Dose: _____ Route: _____

Lab Orders

Required: EBV Seropositive, Neg TB test within 6 months
 Other: _____

Nulojix (belatacept) (Select one):

Initial Dosing: 10 mg/kg IV Day 1, Day 5 end of week 2 and week 4 after transplantation, end of weeks 8 and 12 after transplantation
 Maintenance Dosing: 5 mg/kg at end of week 16 after transplantation, then every 4 weeks (+/-3 days)
 Crossover Dosing: 5 mg/kg on days 1, 15, 29, 43 and 57 followed by 5 mg/kg every 4 weeks

Weight to be used for dosing calculation: _____ kg

Dose based on actual body weight of patient at time of transplant per PI. Dose should be modified if there is a change in body weight of greater than 10%. Dose rounded to nearest 12.5 mg.

Post Treatment Observations: The patient is required to stay for 30 minutes following the first administration.

Refills: zero 6 months 12 months _____ (Prescription valid for one year, unless otherwise indicated)

Special Instructions: _____

PROVIDER INFORMATION

Provider Full Name: _____ Provider NPI #: _____ Specialty: _____

Practice Address: _____ City: _____ State: _____ Zip Code: _____

Contact Name: _____ Phone: _____ Fax: _____ Email: _____

Provider Name (Print)

Provider Signature

Date