

OmvoH (mirikizumab-mrkz)

Provider Order Form rev. 1/12/2026



PATIENT INFORMATION

Referral Status: ☐ New Referral ☐ Updated Order ☐ Order Renewal

Patient Full Name: _____ DOB: _____ Phone: _____ Gender: ☐ M ☐ F ☐ Other

Email Address: _____ Address: _____ Weight (lbs/kg): _____ Height (in): _____

☐ NKDA Allergies: _____ Existing prior authorization? ☐ Yes, (Send a copy) ☐ No (AIC will process)

Patient Status: ☐ New to Therapy ☐ Continuing Therapy Last Treatment Date: _____ Next Due Date: _____

Patient Preferred Location: _____

DIAGNOSIS & CLINICAL INFORMATION

ICD 10-Code & Description (Provide full completed code)

☐ K51.00 Ulcerative (chronic) pancolitis without complications

☐ Other: _____

☐ K51.90 Ulcerative colitis, unspecified without complications

REQUIRED DOCUMENTATION: Please include insurance card (front & back), all patient demographics, history & physicals, medication lists, recent lab results (CBC, CMP, TB, Hep B panel depending on medication), signed prescription order and recent visit notes.

Confirm that these and the required lab orders have been sent to American Infusion Care and necessary parties.

PRESCRIPTION INFORMATION

Nursing: Provide nursing care per American Infusion Care - Specialty Infusions protocols, including reaction management and post-procedure observation

Pre-Medications

☐ Acetaminophen (Tylenol) ☐ 500mg ☐ 650mg ☐ 1000mgPO

☐ Cetirizine (Zyrtec) 10mgPO

☐ Loratadine (Claritin) 10mgPO

☐ Diphenhydramine (Benadryl) ☐ 25mg ☐ 50mg ☐ PO ☐ IV

☐ Methylprednisolone (Solu-Medrol) ☐ 40mg ☐ 125mg IV

☐ Other: _____ Dose: _____ Route: _____

Lab Orders

Required: Liver function tests/Bilirubin, TB Test Results

☐ Other: _____

OmvoH (mirikizumab-mrkz) (Select one):

Ulcerative Colitis:

☐ Induction Dose: Infuse 300 mg IV over at least 30 minutes at Weeks 0, 4, and 8.

☐ Maintenance Dose: Infuse 200mg subcutaneously at week 12 and every 4 weeks thereafter (given as two consecutive injections of 100mg).

☐ Other: _____

Crohn's Disease:

☐ Induction Dose: Infuse 900 mg IV over at least 90 minutes at Weeks 0, 4, and 8.

☐ Maintenance Dose: Infuse 300mg subcutaneously at week 12 and every 4 weeks thereafter (given as two consecutive injections of 100mg and *200mg in any order).

☐ Other: _____

**The 200mg/2mL prefilled pen and syringe are only to be used for the maintenance treatment of Crohn's Disease.*

Post Treatment Observations: The patient is required to stay for 30 minutes following the first administration.

☐ Refills: ☐ zero ☐ 6 months ☐ 12 months ☐ _____ (Prescription valid for one year, unless otherwise indicated)

Special Instructions: _____

PROVIDER INFORMATION

Provider Full Name: _____ Provider NPI #: _____ Specialty: _____

Practice Address: _____ City: _____ State: _____ Zip Code: _____

Contact Name: _____ Phone: _____ Fax: _____ Email: _____

Provider Name (Print)

Provider Signature

Date

E: Referrals@americaninfusioncare.com

Americaninfusioncare.com

Greater Houston Area F: 832.510.7824 P: 832.800.3213
McAllen F: 956.302.8906 P: 832.800.3213 Plano: F: 214.831.9829 P: 972.865.4454
Harlingen F: 956.341.9687 P: 832.800.3213 Laredo F: 956.306.3715 P: 832.800.3213
Other Locations (Oaklawn, Lancaster, etc): F: 469.305.2361 P: 972.865.4454